## **DECISION TREE FOR** Yes -**INHALATION EXPOSURES** Continue ····· Statements in these brackets {...} are optional THREAT TO RESCUER NON-IRRITANT IF INHALED FLAMMABLE? AND NON-TOXIC? (e.g. "knock-down"?) Eliminate all ignition sources Take precautions to ensure your own safety Get medical advice/attention if you feel before attempting rescue (e.g. wear appropriate if safe to do so. unwell or are concerned. protective equipment, use the buddy system). **CAUSES PULMONARY** FATAL/TOXIC IF RESPIRATORY EDEMA OR INTERFERES SENSITIZER? INHALED? WITH OXYGENATION? Remove source of exposure or move person Remove source of exposure or move Remove source of exposure or move person to fresh air and keep comfortable for breathing. Immediately to fresh air and keep comfortable for person to fresh air and keep call a POISON CENTER/doctor/....\* Specific treatment breathing. {Immediately} call a POISON comfortable for breathing. If CENTER/doctor/....\* Specific treatment (is experiencing respiratory symptoms: (is urgent)....\* If breathing is difficult, trained urgent}....\* If breathing has stopped, trained personnel should administer emergency oxygen if Call a POISON CENTER/doctor/....\* If advised to do so by the POISON CENTER/doctor/....\* personnel should begin rescue breathing or, breathing is difficult, trained personnel should administer emergency oxygen if the heart has stopped, immediately start \*...Refer to supplemental first aid instruction, if cardiopulmonary resuscitation (CPR) or if advised to do so by the POISON automated external defibrillation (AED). immediate administration of antidote or specific CENTER/doctor/.... measures required. \*...Refer to supplemental first aid If appropriate, add "Symptoms of pulmonary edema instruction, if immediate administration of can be delayed up to 48 hours after exposure." antidote or specific measures required. If direct contact during rescue breathing poses a threat to the first aid provider, recommend "Avoid If direct contact during rescue breathing mouth-to-mouth contact by using a barrier device." poses a threat to the first aid provider, recommend "Avoid mouth-to-mouth contact by using a barrier device." ORGAN DAMAGE -HARMFUL IF INHALED, RESPIRATORY ORGAN IRRITANT, OR CAUSES SINGLE EXPOSURE DAMAGE - REPEATED DROWSINESS/DIZZINESS? IF INHALED? EXPOSURE? {Take precautions to ensure your own safety (e.g. {If exposed} {If you feel unwell}{If concerned}: Get medical advice/attention if you feel wear appropriate protective equipment).} {Remove Call a POISON CENTER/doctor/....\* {Specific unwell. source of exposure or move person to fresh air treatment...} and keep comfortable for breathing.} Call a POISON CENTER/doctor/....\*/if you feel unwell. ...Refer to supplemental first aid instruction, if immediate measures required.

IF exposed or concerned: Get medical advice/attention.

CARCINOGEN, TOXIC (A TO REPRODUCTION, MUTAGEN?

\*...specify the appropriate source of emergency medical advice.